

# The jasmonic acid signaling pathway is linked to auxin homeostasis through the modulation of *YUCCA8* and *YUCCA9* gene expression

Mathias Hentrich<sup>1</sup>, Christine Böttcher<sup>1,†</sup>, Petra Dückting<sup>1</sup>, Youfa Cheng<sup>2,‡</sup>, Yunde Zhao<sup>2</sup>, Oliver Berkowitz<sup>3,§</sup>, Josette Masle<sup>3</sup>, Joaquín Medina<sup>4</sup> and Stephan Pollmann<sup>4,\*</sup>

<sup>1</sup>Department of Plant Physiology, Ruhr-University Bochum, Universitätsstraße 150, 44780 Bochum, Germany,

<sup>2</sup>Section of Cell and Developmental Biology, University of California at San Diego, 9500 Gilman Drive, La Jolla 92093-0116, CA, USA,

<sup>3</sup>Research School of Biology, Australian National University, Canberra, ACT 2601, Australia, and

<sup>4</sup>Centro de Biotecnología y Genómica de Plantas (CBGP), Campus de Montegancedo, 28223 Pozuelo de Alarcón (Madrid), Spain

Received 7 November 2012; revised 24 January 2013; accepted 12 February 2013; published online 20 February 2013.

\*For correspondence (e-mail stephan.pollmann@upm.es).

† Present address: CSIRO Plant Industry, Hartley Grove, Urrbrae, SA 5064, Australia.

‡ Present address: Institute of Botany, Chinese Academy of Sciences, No. 20 Nanxincun, Xiangshan, Beijing 100093, China.

§ Present address: School of Biological Sciences and Biotechnology, Murdoch University, South St, Murdoch, WA 6150, and School of Plant Biology and Institute of Agriculture, The University of Western Australia, 35 Stirling Highway, Crawley, Perth, WA 6009, Australia.

## SUMMARY

Interactions between phytohormones play important roles in the regulation of plant growth and development, but knowledge of the networks controlling hormonal relationships, such as between oxylipins and auxins, is just emerging. Here, we report the transcriptional regulation of two *Arabidopsis* *YUCCA* genes, *YUC8* and *YUC9*, by oxylipins. Similar to previously characterized *YUCCA* family members, we show that both *YUC8* and *YUC9* are involved in auxin biosynthesis, as demonstrated by the increased auxin contents and auxin-dependent phenotypes displayed by gain-of-function mutants as well as the significantly decreased indole-3-acetic acid (IAA) levels in *yuc8* and *yuc8/9* knockout lines. Gene expression data obtained by qPCR analysis and microscopic examination of promoter-reporter lines reveal an oxylipin-mediated regulation of *YUC9* expression that is dependent on the COI1 signal transduction pathway. In support of these findings, the roots of the analyzed *yuc* knockout mutants displayed a reduced response to methyl jasmonate (MeJA). The similar response of the *yuc8* and *yuc9* mutants to MeJA in cotyledons and hypocotyls suggests functional overlap of *YUC8* and *YUC9* in aerial tissues, while their function in roots shows some specificity, probably in part related to different spatio-temporal expression patterns of the two genes. These results provide evidence for an intimate functional relationship between oxylipin signaling and auxin homeostasis.

**Keywords:** *YUCCA* genes, auxin, jasmonic acid, hormone signaling, plant hormone interaction, root development, *Arabidopsis*.

## INTRODUCTION

Oxylipins, such as jasmonates and their precursor 12-oxo-phytodienoic acid (OPDA), are pleiotropic plant growth regulators that are widely represented throughout the plant kingdom (Böttcher and Pollmann, 2009). These bioactive signaling molecules regulate various plant developmental processes including anther dehiscence, pollen maturation and root growth. In addition, they are well-known mediators of abiotic and biotic stress responses, e.g. to pathogen attack and herbivory (Wasternack, 2007; Howe and Jander, 2008).

There is mounting evidence of extensive interaction between phytohormones in the regulation of plant growth and development or their adaptation to external stimuli (Wolters and Jürgens, 2009). These interactions are in some cases governed by shared components of signal transduction pathways, but in other cases reflect the modulation that a given hormone exerts on the synthesis or action of another one (Hoffmann *et al.*, 2011). Jasmonates are no exception and numerous interactions of jasmonates with other hormones have been reported (Kazan and

Manners, 2008). Among those, the interaction between jasmonic acid (JA) and auxin is still poorly understood despite growing evidence for significant interactions between these two seemingly antagonistic signaling molecules, in particular in regulating the initiation of adventitious roots (Gutierrez *et al.*, 2012) and coleoptile photo- and gravitropism (Riemann *et al.*, 2003; Gutjahr *et al.*, 2005). In addition, there seems to be an intimate interplay between JA and indole-3-acetic acid (IAA) in the control of lateral root formation (Sun *et al.*, 2009).

Besides shared components in ubiquitin-dependent protein degradation pathways such as AXR1 and AXR6 (Tiryaki and Staswick, 2002; Ren *et al.*, 2005), and shared co-repressors like TPL (Szemenyei *et al.*, 2008; Pauwels *et al.*, 2010), auxin and JA pathways are linked through the action of auxin response factors (ARFs) which regulate the transcription of auxin-dependent genes. In addition to an impaired auxin response, *arf6/arf8* mutants exhibit aberrant flower development, and JA deficiency as well as reduced transcription of JA biosynthetic genes in their flowers (Nagpal *et al.*, 2005). The essential role of JA in the regulation of flower development has been previously documented (Ishiguro *et al.*, 2001). Moreover, auxin is capable of inducing the expression of JA biosynthetic genes (Tiryaki and Staswick, 2002), while JA increases the metabolic flux into the L-tryptophan (Trp) pathway by inducing gene expression of ANTHRANILATE SYNTHASE  $\alpha 1$  (*ASA1*) and ANTHRANILATE SYNTHASE  $\beta 1$  (*ASB1*). Elevated *ASA1* and *ASB1* expression leads to increased production of Trp, a precursor for the synthesis of active auxin, namely IAA, and indole glucosinolate (Dombrecht *et al.*, 2007; Sun *et al.*, 2009). Recently, jasmonates have also been reported to be involved in the regulation of auxin transport through effects on the distribution of the auxin exporter PINFORMED 2 (*PIN2*) (Sun *et al.*, 2011).

Although the intricacies of IAA biosynthesis still need to be fully elucidated, both genetic and biochemical studies suggest that YUCCA enzymes play a role in the production of IAA (Mashiguchi *et al.*, 2011; Stepanova *et al.*, 2011; Won *et al.*, 2011; Dai *et al.*, 2013). In Arabidopsis, the YUCCA gene family consists of 11 members encoding flavin-containing monooxygenase-like proteins. Several YUCCA enzymes, e.g. YUC1–6 and YUC10 and -11, have been implicated in the local production of auxin (Zhao *et al.*, 2001; Woodward *et al.*, 2005; Cheng *et al.*, 2006, 2007; Kim *et al.*, 2007, 2011) and the transcription of the YUCCA genes appears to be strictly spatio-temporally regulated (Zhao, 2008). Knowledge of the transcriptional networks by which the expression of the YUCCA genes is regulated, however, is just emerging.

Recent studies attracted our attention by providing circumstantial evidence for the induction of two YUCCA genes, *YUC8* and *YUC9*, by wounding. For example, induction of *YUC9* expression was described in response to

OPDA treatment both in the *tga2/5/6* triple-mutant and the wild-type background (Mueller *et al.*, 2008). Another dataset from an Arabidopsis cell culture experiment gave evidence for a four-fold rapid and transient increase of *YUC8* gene expression in response to application of methyl jasmonate (MeJA) (Pauwels *et al.*, 2008). These results prompted us to hypothesize that *YUC8* and *YUC9* may play an important role in linking oxylipin signaling and IAA metabolism.


Here, we analyze these two as yet uncharacterized members of the Arabidopsis YUCCA gene family. We demonstrate a role for the two genes in auxin homeostasis and root development as well as in hypocotyl and leaf phenotypes typically associated with changes in IAA levels. We show that both genes participate in the induction of auxin synthesis by MeJA and other oxylipins. Our data provide evidence for partial redundancy between the two genes, but also reveal specific functions with respect to their spatio-temporal expression patterns as well as their involvement in the interplay of auxin and oxylipins.

## RESULTS

### YUC8 and YUC9 contribute to auxin production *in vivo*

As a first step to investigate the function of *YUC8* and *YUC9*, we generated stable *35S::YUC8* and *35S::YUC9* overexpression lines (*YUC8ox*, *YUC9ox*). Several independent homozygous *YUC8ox* and *YUC9ox* lines were recovered. All displayed a phenotype similar to other auxin overproducers, e.g. *yucca*, *CYP79B2ox* or *rooty* (Zhao *et al.*, 2002), with elongated hypocotyls, epinastic cotyledons, reduced primary root lengths and narrow elongated leaf blades and petioles (Figure 1a). Hypocotyl length in *YUC8ox* and *YUC9ox* was 1.6- and two-fold greater, respectively, than in wild type, similar to hypocotyl elongation earlier observed in *sur1/rooty* or *sur2* mutants, showing a two- to three-fold increase (Boerjan *et al.*, 1995; King *et al.*, 1995). It is noteworthy, however, that free IAA levels in both *YUC8ox* and *YUC9ox* (Figure 1b), as well as in other previously characterized YUCCA overexpressing lines (Zhao *et al.*, 2001) are only 1.5- to two-fold higher than those in the wild type. This contrasts with the five- to 20-fold increase of IAA contents in *sur1/rooty* and *sur2* (King *et al.*, 1995; Sugawara *et al.*, 2009). This may emphasize the importance of the spatio-temporal distribution of IAA-overproduction, constitutive and unspecific in the YUCox lines due to the use of a 35S promoter to drive overexpression, but most probably tissue/cell specific in the *superroot* knockout mutants.

Like *35S::YUC6* transgenic plants (Kim *et al.*, 2011), adult *YUC8ox* and *YUC9ox* plants showed considerable longevity, with some lines reaching heights of more than 1 m (Figure 1c). To test the hypothesis that elevated IAA levels were the cause of the aberrant phenotypes associated with *YUC8/9* overexpression, auxin levels were measured in a



**Figure 1.** Phenotype of *YUC8* and *YUC9* overexpression lines.

(a) One-week-old and 6-week-old plants of wild type (WT) as well as *YUC8ox* and *YUC9ox* lines. Scale bars, 1 mm on the left, 5 mm on the right. (b) Elevated levels of indole-3-acetic acid (IAA) in *YUC8*- and *YUC9*-overexpressing lines. The IAA levels were determined by gas chromatography-mass spectrometry analysis. For each construct, three independent lines were analyzed and compared with WT. The IAA levels are normalized to 100% in WT. Means are given with their SE ( $n = 6$ ). Similar results were obtained in two independent experiments. Different letters indicate significant differences between means ( $P < 0.05$ ). (c) Arabidopsis overexpressing *YUC8* exhibit strongly enhanced shoot growth and stem elongation. Numbers in the figure indicate plant heights in centimeters. Similar results were obtained for 35S::*YUC9* lines, showing maximum heights of 120 cm (data not shown).

number of independent transgenic lines by gas chromatography–mass spectrometry. In agreement with previous reports on auxin contents in other *YUCCA*-overexpressing lines (Zhao *et al.*, 2001), endogenous auxin levels in our lines were increased by 1.5- to two-fold when compared to wild-type levels (Figure 1b). Consistent with the assumption of a role of *YUCCA* genes in Trp-dependent auxin synthesis (Zhao *et al.*, 2001; Mashiguchi *et al.*, 2011; Stepanova *et al.*, 2011; Won *et al.*, 2011), the examined lines showed an increased resistance towards toxic 5-methyl-tryptophan (5-mT) (Figure S1). Even though 5-mT is ultimately lethal to wild-type and *YUCCA*-overexpressing plants by inhibiting Trp formation and blocking the function of proteins in which this amino acid is incorporated (Hull *et al.*, 2000), it seems as if part of the 5-mT is shunted into non-toxic 5-methyl IAA thereby increasing the resistance of *YUC8ox* and *YUC9ox*.

We also used the 35S::*YUC8* and 35S::*YUC9* overexpression vectors to perform transient assays on *Nicotiana*


*benthamiana* leaves. Within 18 h post-infiltration, leaves infiltrated with either construct showed strong curling compared with control leaves infiltrated with *Agrobacterium tumefaciens* harboring an empty vector (Figure S2a, b). Inspection of leaf epidermis revealed that expression of both *YUC8* and *YUC9* caused a 2–2.5-fold increase in pavement cell size (Figure S2c–e). These observations are in agreement with previous studies on auxin overproducers showing similar leaf curling (Zhao *et al.*, 2001; Zhang *et al.*, 2007).

To confirm the results obtained in our *YUC8* and *YUC9* overexpression experiments, we analyzed IAA levels in 7-day-old *yuc8* and *yuc9* single- and *yuc8/yuc9* double-knockout mutants. Relative to the wild type, two out of three mutants showed significantly reduced IAA contents (Figure 2a). The most severe reduction was detected in *yuc8/yuc9* double-knockout seedlings with an IAA level approximately 50% of that in the wild type compared to 70% for the *yuc8* single mutant. The impact of *YUC9* loss-of-function on the endogenous IAA pool was not statistically significant, with a decrease of only 5–10% compared with wild-type levels. All mutants showed slightly reduced hypocotyl (75%) and petiole (83%) lengths (Figure 2b,c), in line with observations in other mutants with low auxin contents such as the *ilr1/iar3* double- and the *ilr1/iar3/ill2* triple-knockout mutants or the *sav3-2* mutant (Rampey *et al.*, 2004; Tao *et al.*, 2008). Taken together, these results indicate that *YUC8* and *YUC9* both contribute to auxin synthesis in Arabidopsis. In addition, the additive effects on the IAA levels detected in the mutant seedlings under normal conditions (Figure 2a) reveal that *YUC8* and *YUC9* seemingly act in parallel to maintain auxin homeostasis.

### *YUC8* and *YUC9* have distinct expression patterns

To examine the degree of redundancy between the two genes, we analyzed tissue-specific expression levels by quantitative (q)RT-PCR (Figure 3). *YUC8* expression was highest in very young 3- and 7-day-old seedlings, thereafter declining. In adult 6–8-week-old plants, *YUC8* transcript levels were highest in young sink leaves, lower in mature source leaves, roots and flower buds, and hardly detectable in stems (Figure 3a). *YUC9* gene expression was generally one order of magnitude lower than *YUC8* expression (Figure 3b). It was also developmentally regulated in young seedlings but with a peak 7 days after germination. In adult plants, *YUC9* was predominantly expressed in roots and at a much lower level in sink leaves, while being undetectable in floral tissue. These patterns are consistent with microarray data deposited in public databases, e.g. the Genevestigator database (Hruz *et al.*, 2008), and indicate marked differences in the spatio-temporal expression patterns of the two genes.

We next generated *pYUC8*::GUS and *pYUC9*::GUS reporter lines allowing for finer resolution in the examination of




**Figure 2.** Endogenous auxin contents and phenotype of *yuc8* and *yuc9* single- and double-knockout mutants.

(a) Indole-3-acetic acid (IAA) content in 7-day-old seedlings. Auxin content was quantified by gas chromatography-mass spectrometry analysis. Means are given with their SE ( $n = 6$ ). Similar results were obtained in three independent experiments. Different letters indicate significant differences between means ( $P < 0.05$ ). Fw, fresh weight.

(b), (c) Seedlings grown for 7 days on vertical half-strength MS plates were analyzed for (b) their petiole and (c) their hypocotyl length. Both petioles and hypocotyls are significantly shorter in the mutants. Given are means with their SE ( $n = 8-10$  seedlings). Different letters indicate significant differences between means ( $P < 0.05$ ).

cellular and tissue specificity in *YUC8* and *YUC9* expression. Within 2 days of germination, expression of both *YUC8* and *YUC9* was detectable in primary root tips, and within the next 2 days became visible in the root



**Figure 3.** Tissue specific expression of

(a) *YUC8* and

(b) *YUC9* in wild type. Transcript abundance values are given relative to the geometric means of *APT1* and *UBQ10* transcripts. Error bars are given  $\pm$ SE.

vasculature (Figure S3). In older plants, GUS reporter activity was absent from leaves (*YUC8*) or confined to the very margin of sink leaves (*YUC9*, Figure S4h). The two genes were expressed in reproductive organs but with quite distinct patterns. *YUC8* showed high specific expression in flower buds (Figure S4c). Closer examination of flowers revealed *YUC8* promoter activity in the young anthers of immature flowers (Figure S4i) and in the vascular tissues of sepals and petals of mature flowers as well as in anther filaments (Figure S4i-m) and the funiculus and integuments of young ovules (Figure S5b-d). These results extend the previously described pattern of *YUC8* promoter activity (Rawat *et al.*, 2009) and suggest a role for *YUC8* in reproductive development. By contrast, there was no detectable *YUC9* promoter activity in floral tissues, but transient expression in the suspensor of the embryo, at the 8- and 16-cell stage (Figure S5a,e-g). As suspensor cells are known to synthesize IAA (Kawashima and Goldberg, 2010), the expression pattern of *YUC9* suggests a role for *YUC9* in supply of IAA during early embryo development. Overall, and consistent with our qRT-PCR data, these results confirm tissue specificity in *YUC8* and *YUC9* gene

expression. In order to compare the subcellular localization of the encoded proteins, both genes were fused to the green fluorescent protein (*GFP*) reporter gene. The resulting constructs were then used to transiently transfect Arabidopsis pavement cells by ballistic gene transfer. Confocal laser scanning microscopy revealed an exclusive cytoplasmic localization of the fusion proteins (Figure S6).

#### **YUC8 and YUC9 contribute to MeJA-induced auxin synthesis at the organ level**

Previous studies reported that treatment with either MeJA or coronatine promotes the production of auxin (Uppalapati *et al.*, 2005; Dombrecht *et al.*, 2007; Sun *et al.*, 2009). Hence, we next analyzed auxin levels and responsiveness to MeJA application in the wild type and *yuc8* and *yuc9* single- and double-knockout mutants. Seven-day-old seedlings were treated for 2 h with either 50  $\mu\text{M}$  MeJA or a control solution (0.5% methanol, v/v) and then dissected into cotyledons, hypocotyls and roots for separate analysis of IAA contents. Endogenous IAA levels in control mock-treated seedlings were affected by *YUC* loss-of-function to varying degrees depending on the organ considered, with no effect in cotyledons, a decrease in hypocotyls in *yuc9* and more so in *yuc8*, and for roots a decrease in the *yuc8* mutant only (Figure S7). Combining the two mutations had no effect in cotyledons, but data for the double mutant suggested some additive effects of the *yuc8* and *yuc9* mutations in hypocotyls while in roots IAA levels in the double mutant mirrored those measured in *yuc8*. Application of MeJA consistently caused an elevation of IAA contents (Figure 4a), but the extent of this effect was organ- and line-dependent. While the double mutant showed reduced responsiveness to MeJA in all organs, single mutations did not significantly affect responsiveness. These results indicate a role for both YUC8 and YUC9 in overall auxin homeostasis and MeJA-induced IAA production.

#### **The root response to MeJA treatment is impaired in *yuc8* and *yuc9***


Wild-type seedlings respond to jasmonate treatment by a reduction of primary root growth (Figure 4b), consistent with earlier reports (Staswick *et al.*, 1992). This MeJA-induced inhibition of root elongation was partially suppressed in all mutant lines (Figure 4b), most significantly so in the *yuc8/yuc9* double mutant. Primary roots of MeJA-treated seedlings of this mutant were 53% shorter than corresponding untreated *yuc8/yuc9* mutants, while treated *yuc8* and *yuc9* mutants showed 62% shorter primary roots. Under these same conditions primary roots in MeJA-treated wild-type seedlings were 82% shorter relative to the untreated wild-type plants. These gradual effects are in agreement with the differential reductions in MeJA-induced IAA synthesis in these mutants (Figure 4a).

Methyl jasmonate treatment also caused a considerable reduction in the number of lateral roots, in all lines (Figure 4c). In the wild type this reduction was of relatively smaller magnitude than in the *yuc8* and *yuc9* single and double mutants. While in MeJA-treated wild-type seedlings the lateral root number is decreased to 48% of that in the untreated control, the reduction was more pronounced in *yuc9* and *yuc8/yuc9* mutants, with a decrease to 27% and 30%, respectively, relative to the corresponding untreated samples. However, when combining the observation of inhibition of primary root growth with those on lateral root numbers, considering lateral root density as the index figure, we were able to detect that MeJA treatment did not result in a diminished lateral root density, which is probably due to the pronounced inhibition of primary root growth (Figure 4d). While wild-type seedlings responded to MeJA application with a nearly two-fold increase in lateral root density, this promoting effect was partly suppressed by the *yuc8* mutation and completely abolished by the *yuc9* mutation. In both the *yuc9* and *yuc8/yuc9* double mutants, MeJA no longer enhanced lateral root formation: lateral root density was similar on treated and untreated roots. These results show that *YUC9* and possibly *YUC8* are involved in mediating the promoting effect of MeJA on lateral root formation.

The different responses to MeJA of the *yuc8* and *yuc9* mutants prompted us to more closely examine the expression patterns of the two genes in roots using our GUS reporter lines. Both lines were consistently expressed in root tips (Figure 5a–d), *YUC8* promoter activity was visible early during lateral root formation at the sites of primordia initiation and was very high in the vasculature of the young emerged lateral roots (Figure 5e). *YUC9* expression was detectable even earlier, but was more diffuse in young lateral roots and also expressed in the vasculature of subtending roots (Figure 5f). These not totally overlapping expression patterns may suggest distinct roles of *YUC8* and *YUC9* in the local auxin supply through which the growth of lateral roots may be supported. Overall, these data suggest that YUC8 and YUC9 promote lateral root formation, possibly in a synergistic manner.

#### **Induction of *YUC9* expression by oxylipins is COI1 dependent**

Our findings here of an increase in IAA levels after MeJA treatment as well as published gene expression datasets (Mueller *et al.*, 2008; Pauwels *et al.*, 2008) indicated oxylipin-dependent changes of *YUC8* and *YUC9* gene expression and/or activity. We therefore investigated the hypothesis that the reduced responsiveness of the *yuc8* and *yuc9* mutants to MeJA may reflect a role of oxylipins on the transcriptional regulation of *YUC8* and *YUC9*. To address this, we measured *YUC8* and *YUC9* transcript abundance by qRT-PCR (Figure 6) following application of




**Figure 4.** Comparison of methyl jasmonate (MeJA)-induced auxin formation as well as root phenotypes in wild type (WT) and both *yuc8* and *yuc9* single and double mutants.

(a) Auxin accumulation after MeJA treatment is given by the difference of indole-3-acetic acid (IAA) levels in mock-treated samples and those in MeJA-treated ones. The corresponding absolute amounts are given in Figure S7. After the treatment seedlings were dissected to separate cotyledons, hypocotyls and roots. All organs were immediately frozen in liquid nitrogen. Means are given  $\pm$  SE ( $n = 6$ ). Similar results were obtained in three independent experiments.

(b) Analysis of WT, *coi1*, *yuc8*, *yuc9* and *yuc8/yuc9* mutant primary root lengths. Without supplementation the primary root length of the compared lines does not significantly differ. When grown on media containing 20  $\mu$ M MeJA, the examined *yuc* mutant roots are significantly longer than the WT ones; however, they are considerably shorter than *coi1* primary roots. Means are given with their SE ( $n = 8$ –10 seedlings).

(c) Relative to WT roots, the roots of *yuc8* and *yuc9* single and double mutants show a significantly reduced number of lateral roots. Moreover, lateral root density given in number of lateral roots calculated per centimeter of primary root is lower (d). For comparison, root systems of seedlings grown on half-strength MS medium or medium supplemented with MeJA were analyzed. Illustrated are means  $\pm$  SE ( $n = 15$ –20). Similar results were obtained in three independent experiments. Mean values within a graph are significantly different ( $P < 0.05$ ) where superscript letters differ.



**Figure 5.** Histochemical staining of root tips and emerging lateral roots in transgenic promoter reporter lines.

Depicted are GUS-stained primary root tips of

(a) *pYUC8::GUS* and

(c) *pYUC9::GUS* at 1, 2, 4, 7, 10 and 14 days after germination.

(b) *YUC8* and


(d) *YUC9* promoter activity in the tips of lateral roots at 8, 12 and 14 days after emergence from the epidermis. Scale bars, 50  $\mu$ m (a–d). b-Glu-curonidase activity in

(e) *pYUC8::GUS* and

(f) *pYUC9::GUS* reporter lines during lateral root formation. Plants were grown in sterile conditions on plates and then stained for reporter activity. Scale bars: 100  $\mu$ m (e, f).

MeJA, OPDA or coronatine, a bacteriotoxin that mimics the bioactive jasmonate JA-Ile (Fonseca *et al.*, 2009). In wild-type seedlings, all treatments resulted in a statistically significant reduction of *YUC8* transcript levels within 30 min, followed by a progressive recovery within the next 3 h (Figure 6a). This pattern is in line with published expression data on transcriptional responses mediated by jasmonates (Mandaokar *et al.*, 2006), from which one can extract a similar kinetic of *YUC8* transcripts. The *YUC8* induction by MeJA previously observed in cell culture (Pauwels *et al.*, 2008) may result from a transient and more general stress response and not from a specific oxylipin

induction as suggested by the recent report that protoplasting of root tissue leads to a 10-fold increase in *YUC8* expression (Dinneny *et al.*, 2008). In contrast to *YUC8*, *YUC9* transcription was not repressed but rather induced by oxylipins (Figure 6a), with a seven-fold increase in transcript levels after 120 min of OPDA treatment while MeJA and coronatine caused an even stronger induction (11- and 12-fold increase in transcript abundance 240-min post-treatment, respectively). *YUC9* induction by MeJA is consistent with the increases in IAA levels and alterations of




**Figure 6.** Oxylin-mediated induction of *YUCCA* gene expression. Quantitative real-time reverse transcriptase PCR for *YUC8* and *YUC9* gene expression in wild type. (a), and both *yuc8* and *yuc9* (b), as well as *coi1* (c) mutant seedlings after treatment under variable conditions. Seedlings were either treated with a 50  $\mu$ M methyl jasmonate (MeJA), 50  $\mu$ M 12-oxo-phytodienoic acid (OPDA), 10  $\mu$ M coronatine or control solution (half-strength MS medium, 0.5% methanol v/v). The duration of induction varied between 30 and 240 min (a) and 2 h (b, c). Additionally, a long-term treatment with 20  $\mu$ M MeJA in plates was conducted (b). Then, seedlings were harvested, total RNA extracted, and transcript levels examined by quantitative RT-PCR. Relative expression levels of the genes are normalized to levels in mock controls or untreated control samples and represent the average of three biological replicates, with error bars denoting the standard error. Student's *t*-test; \*\**P* < 0.01. Mean values within a graph are significantly different (*P* < 0.05) where superscript letters differ.

root development shown in Figure 4. The transient down-regulation of *YUC8* was, however, unexpected given the apparent synergy of the two genes in increasing respon-

siveness to oxylin. To further examine the interaction between *YUC8* and *YUC9*, we asked whether a loss of either gene function might be compensated for by an up-regulation of the other one; therefore we analyzed *YUC8/9* expression in the corresponding knockout background (Figure 6b). Relative to untreated seedlings *YUC8* and *YUC9* transcript levels were significantly more abundant after short-term MeJA treatments, which may also explain why *yuc8* mutants show oxylin-related phenotypic defects. These experiments indicate that the two genes can, at least partially, compensate for a loss of function of the other and that both genes are functionally linked. This highlights that in addition to the already reported *YUC8/9* induction by high temperatures (Stavang *et al.*, 2009), the transcriptional regulation of both genes can be considerably affected by MeJA.

Jasmonic acid signal transduction proceeds via the COI1 signal transduction pathway (Feys *et al.*, 1994; Browse, 2009; Chini *et al.*, 2009). The root phenotype of *yuc8* and *yuc9* mutants grown on MeJA-containing media is intermediate between that of wild type and the jasmonate-insensitive *coi1* mutant (Figure 4b). The *yuc8* and *yuc9* single and double mutants, however, retain a residual sensitivity to MeJA, as shown by our phenotypic analyses and quantification of IAA levels, even though their responsiveness is significantly reduced (Figure 4b). To investigate whether oxylin-dependent *YUCCA* expression is also governed by COI1, we treated the *coi1* mutant with the various oxylin and examined changes in *YUC8* and *YUC9* expression (Figure 6c). When compared with responses obtained 2 h after induction of wild-type plants, transcriptional responses to oxylin treatment appeared almost entirely suppressed in the *coi1* background. This suggests that the tested oxylin most likely regulate *YUC8* and *YUC9* transcription through a COI1-dependent pathway (Figure 6c).

In agreement with the qRT-PCR results, the *pYUC9::GUS* reporter showed substantial induction in primary roots transiently treated with 50  $\mu$ M MeJA, while no significant change in *YUC8* promoter activity could be detected (Figure 7b,c). When plants were grown continuously on plates containing MeJA, activation of the *pYUC8::GUS* reporter was detectable but at an obviously lower level than that of the *pYUC9::GUS* reporter (Figure S8). Local application of MeJA on leaves also activated the *YUC9* promoter (Figure 7a). We next examined whether the oxylin levels produced in response to wounding (Glauser *et al.*, 2009) were sufficient to trigger *YUC8* and *YUC9* promoter activation. *YUC9* expression responded strongly, similar to the wound-inducible *ALLENE OXIDE SYNTHASE* promoter (Kubigsteltig *et al.*, 1999), while there was no detectable reporter activity in the *pYUC8::GUS* lines (Figure 7d). Overall, these data indicate that *YUC9* but not *YUC8* expression is up-regulated by jasmonates through a COI1-dependent pathway.



**Figure 7.** Induction of *YUC* gene expression and GUS staining of *pYUC8*:GUS and *pYUC9*:GUS promoter reporter lines after treatment with methyl jasmonate (MeJA).

(a) *YUC9* promoter activation 2 h after application of 10  $\mu$ l 1 mM MeJA on a single leaf (arrow).

(b), (c) Promoter activity in primary root segments of 7-day-old *pYUC8*:GUS and *pYUC9*:GUS lines 2 h after treatment with either a mock control (b) or with a solution containing 50  $\mu$ M MeJA (c). Scale bars: 1 cm (a–c).

(d) Pattern of GUS activity in 4-week-old sterile grown wild type (WT), AOS-promoter GUS line 4-1 (AOS) (Kubigsteltig *et al.*, 1999), *pYUC8*:GUS (*YUC8*) and *pYUC9*:GUS (*YUC9*) lines 12 h after wounding with sterile forceps (arrows). Scale bars: 20  $\mu$ m (d).

## DISCUSSION

Auxin is an essential plant growth hormone ubiquitous throughout the plant kingdom and well known to contribute to a wide array of physiological processes, acting either alone or in interaction with other phytohormones (Davies, 2010). However, the intricacies of the biosynthesis of the primary plant auxin, IAA, are yet to be fully uncovered. Currently, IAA formation is proposed to mainly proceed through four Trp-dependent pathways (Lehmann *et al.*, 2010; Novák *et al.*, 2012). In this context, YUCCA enzymes have received much attention since their discovery. The molecular function of YUCCA monooxygenases has been disclosed recently (Mashiguchi *et al.*, 2011; Stepanova *et al.*, 2011; Won *et al.*, 2011). They are thought

to play a key role in auxin production, catalyzing the conversion of indole-3-pyruvate to IAA. YUCCA monooxygenases have been extensively studied, not only in *Arabidopsis* but also in tomato, maize, petunia and rice (Tobeña-Santamaria *et al.*, 2002; Yamamoto *et al.*, 2007; LeClere *et al.*, 2010; Expósito-Rodríguez *et al.*, 2011), but despite intensive work their transcriptional regulation is still not entirely clear. Thus far, from the 11 YUCCA homologues in *Arabidopsis* eight have been studied in more detail (Zhao *et al.*, 2001; Woodward *et al.*, 2005; Cheng *et al.*, 2006, 2007; Kim *et al.*, 2007, 2011). At present, only the functions of *YUC7–9* have not been settled. The results described here give evidence that, as other YUCCAs, both *YUC8* and *YUC9* also participate in auxin production in *Arabidopsis*. In addition, detailed analysis of *YUC8* and *YUC9* gene expression during the course of plant development and in response to oxylipins demonstrates partial functional overlap between the two genes but also specificity based on their different spatio-temporal expression patterns.

Phenotypically, *yuc8* and *yuc9* single and double mutants grown on MS media do not substantially differ from the wild type, except for slightly shorter hypocotyls and petioles as well as a slightly reduced number of lateral roots (Figures 2 and 4). The root phenotype of the *yuc8* and *yuc9* mutants and the localized GUS staining observed in our reporter lines (Figure 5) imply an involvement of *YUC8* and *YUC9* in lateral root development.

Our data also give evidence for a direct link between oxylipin signaling and auxin biosynthesis. The examined oxylipins seem to transiently repress the expression of *YUC8*, although *YUC8* expression in the *yuc9* background appears to be increased after MeJA treatment (Figure 6a,b). Also, the *YUC8* promoter becomes activated when seedlings are raised on plates containing MeJA (Figure S8). Similarly to the increased *YUC8* response in *yuc9* mutants, induction of *YUC9* expression is more pronounced in *yuc8* seedlings, suggesting that the two genes are functionally linked and that they can, to a certain extent, compensate for each other (Figure 6a,b). The regulatory effect of oxylipins both on *YUC8* and *YUC9* expression is, however, abrogated in the *coi1* background, pointing to an involvement of the COI1-dependent signaling pathway in the regulation of the two genes (Figure 6a,c).

A limited number of previous studies have already provided evidence for a tight connection between JA and auxin homeostasis as well as for an involvement of the jasmonate signaling machinery in the regulation of auxin biosynthesis (for review see, Hoffmann *et al.*, 2011). Those studies revealed a moderate impact of MeJA on *YUC2* expression (Sun *et al.*, 2009), mainly focusing on JA-mediated processes upstream of the auxin biosynthetic pathways (i.e. Trp production), auxin transport or auxin responsiveness (Riemann *et al.*, 2003; Gutjahr *et al.*, 2005;

Dombrecht *et al.*, 2007; Sun *et al.*, 2011). The present results on the regulation of *YUC8* and *YUC9* provide novel insight into the intimate relationship between jasmonate signaling and auxin biosynthesis, pointing out that JA not only triggers substrate production for subsequent auxin synthesis (Sun *et al.*, 2009) but also induces the formation of IAA by the direct induction of IAA synthesis-related genes.

The *asa1-1* mutation appears to interfere with the enhancing effect of MeJA on lateral root formation, which is completely abolished upon treatment with more than 20  $\mu\text{M}$  MeJA (Sun *et al.*, 2009). Under these same conditions (20  $\mu\text{M}$  MeJA), the number of lateral roots in *yuc8* and *yuc9* mutants is also substantially reduced. This effect is most significant in the *yuc8/yuc9* double mutation (Figure 4d), but unlike in *asa1-1*, where lateral root development is totally suppressed, *yuc8/yuc9* mutants still exhibit lateral roots (70% of the number of emerged lateral roots compared to untreated *yuc8/yuc9* seedlings). This indicates that *YUC8* and *YUC9* are probably not the only components of auxin formation downstream of *ASA1*, linking jasmonate signaling to auxin biosynthesis.

In this context, it appears noteworthy that we did not observe an increase in lateral root number upon MeJA treatment as point out previously (Sun *et al.*, 2009). Rather, we detected a substantial decrease (52%) in the number of lateral roots when comparing untreated and 20  $\mu\text{M}$  MeJA-treated wild-type seedlings. This discrepancy may be explained by the fact that Sun *et al.*, (2009) included lateral root primordia in their assessment, while we focused only on lateral roots that had already emerged. On the other hand, differences in growth conditions, e.g. sugar content in the medium, could be responsible for this inconsistency.

While the *asa1-1* mutation does not considerably impact on the MeJA-triggered suppression of primary root elongation, *yuc8* and *yuc9* mutations attenuate the inhibitory effect of MeJA on primary root elongation. This finding extends the previous observations made by Sun *et al.*, (2009). Collectively, these results highlight that application of MeJA triggers an increase in IAA levels by inducing both the production of the major auxin precursor, L-Trp, and auxin synthesis.

Expression of *YUC9* also responds to wounding and MeJA treatment in a COI1-dependent manner (Figures 6a and 7). Moreover, the *yuc8* and *yuc9* knockouts show a root phenotype that is intermediate between the wild type and the JA-insensitive *coi1* mutant, with both *YUCCA* gene products seemingly forming a functional unit acting synergistically on root development and root growth responses. These observations suggest that reduced primary root elongation in response to MeJA treatment is not solely due to primary effects of JA, e.g. through inhibition of mitosis (Zhang and Turner, 2008). Rather, MeJA-mediated

primary root growth inhibition also involves the induction of auxin synthesis and the subsequent action of this additionally formed plant hormone.

As pointed out before, the *asa1-1* mutation does not result in altered primary root elongation responses to MeJA when compared with wild-type plants. Thus, the inhibitory effect of MeJA on root elongation does not seem to depend on *ASA1*. By contrast, the examined combined *yuc8* and *yuc9* mutations do significantly suppress the inhibitory effect of MeJA on primary root elongation (primary roots of MeJA-treated *yuc8/yuc9* mutants appeared 2.9-fold longer than the roots of equally treated wild-type plants, *P*-value = 0.0077) (Figure 4b). Both *ASA1* and the *YUCCA* genes affect IAA levels, but *ASA1* might play a more general role and act on the formation of L-Trp, a primary building block in protein synthesis. Data deposited in public databases (Genevestigator) show a medium to strong *ASA1* expression in nearly all tissues and organs. In contrast, all *YUCCA* genes examined so far are characterized by a very tight spatio-temporal regulation of their expression. This more specific expression pattern may be the reason why *yuc8* and *yuc9* mutations alter the inhibitory effect of MeJA on primary root elongation.

In summary, we propose a modified model in which the observed root growth responses are collectively mediated by the action of JA and IAA. Besides its primary functions, we demonstrate that JA induces the overexpression of *YUC9*. Together with the action of other induced enzymes, the increased YUC levels cause transient IAA overproduction. In fact, this is a good example of an indirect mode of interaction (Hoffmann *et al.*, 2011). In turn, the increased IAA levels result in altered lateral root development and local YUC action seems to also contribute to the inhibition of primary root elongation by MeJA-induced IAA production. As phytohormones act in a dose-dependent manner, with elevated auxin concentrations exerting inhibitory effects on growth (Thimann, 1938), the increased IAA content may reinforce the growth inhibiting effect of jasmonate. A transient, local increase in IAA levels may lead to local growth inhibition and also override positional auxin signals, for example auxin gradients or local auxin maxima. In this context, it is interesting that jasmonates have been reported to interfere with auxin polar transport in a concentration-dependent manner (Sun *et al.*, 2009, 2011).

The relationships between JA signaling and IAA production described here might provide a further regulatory mechanism for the control of the responses of the plant to biotrophic or necrotrophic pathogens. The defense against these two types of pathogens is generally governed by an antagonistic relationship between salicylic acid (SA, induced by biotrophs) and jasmonic acid/ethylene (JA/ET, induced by necrotrophs and herbivores). Several studies have, however, already pointed to an interconnection

between auxin and SA- or JA/ET-dependent pathogen responses (Chen *et al.*, 2007; Truman *et al.*, 2010). As the complex networks of phytohormone interaction are slowly deciphered, further insights on the sophisticated cellular processes that are affected by YUCCA enzymes will be revealed.

## EXPERIMENTAL PROCEDURES

### Plant material and growth conditions

In this study, we analyzed the *Arabidopsis thaliana* ecotype Col-0, the mutants *coi1* (Feys *et al.*, 1994), *yuc8* and *yuc9* (Tao *et al.*, 2008), and the promoter fusions *35S::GUS* (Biesgen and Weiler, 1999) as well as *pAOS::GUS* (Kubigsteltig *et al.*, 1999). Seeds were germinated on solidified half-strength MS ( $\frac{1}{2}$ MS) medium supplemented with 1% sucrose in a growth chamber under controlled conditions ( $110 \mu\text{E}\cdot\text{m}^{-2}\cdot\text{sec}^{-1}$  photosynthetically active radiation, supplied by standard white fluorescent tubes; 8 h light at 24°C, 16 h darkness at 20°C).

### Oxylipin treatment

Oxylipin treatment prior to the quantification of free IAA levels and RNA extraction was carried out as follows. If not stated otherwise, seedlings were grown on  $\frac{1}{2}$ MS medium (1% sucrose) for 7 days. Next, they were transferred to either 50  $\mu\text{M}$  MeJA solution (in liquid  $\frac{1}{2}$ MS medium containing 0.5% methanol v/v) or a control solution (liquid  $\frac{1}{2}$ MS medium, 0.5% methanol v/v) and incubated for the indicated period of time. In case of long-term treatments, i.e. 7 days, seedlings were grown on  $\frac{1}{2}$ MS medium (1% sucrose) containing 20  $\mu\text{M}$  MeJA and taken directly from the plates. Control plants were grown on similar plates lacking MeJA. For the GUS staining of roots on plates (Figure 7b,c), seedlings of the reporter lines were grown on vertical  $\frac{1}{2}$ MS medium plates supplemented with 1% sucrose. To trigger responses, 5 ml of either a mock or a 50  $\mu\text{M}$  MeJA solution (see above) was administered to the plates. After the indicated period of time the plants were washed with  $\frac{1}{2}$ MS solution before they were stained for  $\beta$ -glucuronidase activity as explained below.

### RT-PCR

Reverse transcription was performed on total RNA isolated from wild-type plants as well as *yuc8*, *yuc9* and *coi1* mutant lines. RNA was prepared from 100 mg plant tissue by using TRIzol reagent (Invitrogen, <http://www.invitrogen.com/>) according to the manufacturer's instructions. The mRNA was further purified using an Oligotex mRNA kit (Qiagen, <http://www.qiagen.com/>). First-strand synthesis was performed according to the supplier's instructions, using Moloney murine leukemia virus reverse transcriptase and oligo(dT)<sub>15</sub> primer (Promega, <http://www.promega.com/>).

### Expression analysis by qRT-PCR

Seedlings were grown, RNA isolated and qRT-PCR carried out as previously described (Lehmann and Pollmann, 2009). Samples were run in triplicate. Relative quantification of expression was calculated after data analysis using the Opticon monitor software (v. 2.02.24, MJ Research, <http://www.bio-rad.com/>) by the comparative  $2^{-\Delta\Delta CT}$  method (Livak and Schmittgen, 2001) with *APT1* and *UBQ10* as reference genes (Czechowski *et al.*, 2005). For validation of reference genes and determination of amplification efficiencies the geNORM and LinRegPCR algorithms, respectively, were used

(Vandesompele *et al.*, 2002; Ramakers *et al.*, 2003) Primer sets were as follows: *APT1* (*At1g27450*), Pri15-Pri16; *UBQ10* (*At4g05320*), Pri17-Pri18; *YUC8*, Pri19-Pri20; *YUC9*, Pri21-Pri22 (primer sequences are given in supporting Table S1).

### GUS staining

All analyzed seedlings were grown on plates as described above. For analysis of the older plants and generative organs, plantlets were transferred into liquid MS medium (1% sucrose) 2 weeks after germination. Plant material was pre-cleared with 90% acetone for 0.5 h and then stained in 50 mM citrate buffer (pH 7) containing 1 mM potassium ferrocyanide, 1 mM potassium ferricyanide, 0.1% Triton X-100 and 1 mM 5-bromo-4-chloro-3-indolyl  $\beta$ -D-glucuronide cyclohexylammonium salt for 14 h at room temperature (22 °C). Thereafter, plant material was dehydrated with an ethanol series (20, 35, 50, 70, 80, 90% v/v). Prior to microscopy, GUS-stained tissues were incubated in Hoyers solution overnight (Liu and Meinke, 1998).

### Microscopy

*Arabidopsis* leaves were transformed by particle bombardment and the microscopic analysis of the distribution of GFP-tagged fusion proteins was carried out as previously described (Pollmann *et al.*, 2006). After incubation for 16–18 h, the YUC-GFP expressing cells were analyzed by confocal laser scanning microscopy.

The GUS-stained tissues were examined with a Zeiss Axioskop 100. Images were captured on a Zeiss AxioCam MRc camera (<http://www.zeiss.com/>). Whole seedlings were analyzed with a Leica MZ12 microscope (<http://www.leica.com/>) and digital images were captured with a Nikon D200 camera (<http://www.nikon.com/>).

### Auxin quantification

Whole or dissected seedlings were immediately frozen in liquid N<sub>2</sub>. About 100 mg of tissue was pooled per sample and at least three biological replicates were harvested for each independent experiment. The IAA was quantified on a Varian Saturn 2000 GC-MS/MS system as previously described (Pollmann *et al.*, 2009).

### Statistical analysis

The data were analyzed with one-way ANOVA followed by Tukey's B post-hoc test to allow for comparisons among all means or with Student's *t*-test when two means were compared. Statistical analyses were conducted using PRISM version 5.03 (GraphPad Software, <http://www.graphpad.com/>).

### ACKNOWLEDGEMENTS

The work was supported by DFG grant SFB480-A10, MICINN grant BFU2011-25925, Marie-Curie grant FP7-PEOPLE-CIG-2011-303744, by a grant of the Ruhr-University Bochum (RUB) to promote young academics (SP), and by NIH grant R01GM68631 (YZ). MH was supported by a fellowship from the German National Academic Foundation and a membership of the Research School of the RUB.

### SUPPORTING INFORMATION

Additional Supporting Information may be found in the online version of this article.

**Figure S1.** Growth of YUC8ox and YUC9ox on 5-methyl-tryptophan-containing medium.

**Figure S2.** Transient expression of YUC8 and YUC9 in *Nicotiana benthamiana*.

**Figure S3.** *YUC8* and *YUC9* expression during seedling development.

**Figure S4.** GUS pattern in flowers and leaves.

**Figure S5.** *YUC8* and *YUC9* expression in siliques and ovules.

**Figure S6.** Intracellular localization of *YUC8*- and *YUC9*-GFP fusion proteins.

**Figure S7.** GC-MS analysis of indole-3-acetic acid levels in wild type and *yuc* mutant seedlings.

**Figure S8.** *pYUC8::GUS* and *pYUC9::GUS* lines grown on plates containing methyl jasmonate.

**Methods S1.** Generation of transgenic plants.

**Table S1.** List of primers used in this study.

## REFERENCES

- Biesgen, C. and Weiler, E.W. (1999) Structure and regulation of *OPR1* and *OPR2*, two closely related genes encoding 12-oxophytodieneoic acid-10,11-reductases from *Arabidopsis thaliana*. *Planta*, **208**, 155–165.
- Boerjan, W., Cervera, M.T., Delarue, M., Beeckman, T., Dewitte, W., Bellini, C., Caboche, M., van Onckelen, H., van Montagu, M. and Inze, D. (1995) *superroot*, a recessive mutation in *Arabidopsis*, confers auxin overproduction. *Plant Cell*, **7**, 1405–1419.
- Böttcher, C. and Pollmann, S. (2009) Plant oxylipins: plant responses to 12-oxo-phytodieneoic acid are governed by its specific structural and functional properties. *FEBS J.* **276**, 4693–4704.
- Browse, J. (2009) Jasmonate passes muster: a receptor and targets for the defense hormone. *Annu. Rev. Plant Biol.* **60**, 183–205.
- Chen, Z., Agnew, J.L., Cohen, J.D., He, P., Shan, L., Sheen, J. and Kunkel, B.N. (2007) *Pseudomonas syringae* type III effector AvrRpt2 alters *Arabidopsis thaliana* auxin physiology. *Proc. Natl Acad. Sci. USA*, **104**, 20131–20136.
- Cheng, Y., Dai, X. and Zhao, Y. (2006) Auxin biosynthesis by the YUCCA flavin monooxygenases controls the formation of floral organs and vascular tissues in *Arabidopsis*. *Genes Dev.* **20**, 1790–1799.
- Cheng, Y., Dai, X. and Zhao, Y. (2007) Auxin synthesized by the YUCCA flavin monooxygenases is essential for embryogenesis and leaf formation in *Arabidopsis*. *Plant Cell*, **19**, 2430–2439.
- Chini, A., Boter, M. and Solano, R. (2009) Plant oxylipins: COI1/JAZs/ MYC2 as the core jasmonic acid-signalling module. *FEBS J.* **276**, 4682–4692.
- Czechowski, T., Stitt, M., Altmann, T., Udvardi, M.K. and Scheible, W.R. (2005) Genome-wide identification and testing of superior reference genes for transcript normalization in *Arabidopsis*. *Plant Physiol.* **139**, 5–17.
- Dai, X., Mashiguchi, K., Chen, Q., Kasahara, H., Kamiya, Y., Ojha, S., Dubois, J., Ballou, D. and Zhao, Y. (2013) The Biochemical Mechanism of Auxin Biosynthesis by an *Arabidopsis* YUCCA Flavine-containing Monooxygenase. *J. Biol. Chem.* **288**, 1448–1457.
- Davies, P.J. (2010) *Plant Hormones. Biosynthesis, Signal Transduction, Action!* Revised 3rd edn. Dordrecht, Boston, London: Kluwer Academic Publishers.
- Dinneny, J.R., Long, T.A., Wang, J.Y., Jung, J.W., Mace, D., Pointer, S., Barron, C., Brady, S.M., Schiefelbein, J. and Benfey, P.N. (2008) Cell identity mediates the response of *Arabidopsis* roots to abiotic stress. *Science*, **320**, 942–945.
- Dombrecht, B., Xue, G.P., Sprague, S.J. et al. (2007) MYC2 differentially modulates diverse jasmonate-dependent functions in *Arabidopsis*. *Plant Cell*, **19**, 2225–2245.
- Expósito-Rodríguez, M., Borges, A.A., Borges-Pérez, A. and Pérez, J.A. (2011) Gene structure and spatiotemporal expression profile of tomato genes encoding YUCCA-like flavin monooxygenases: the *ToFZY* gene family. *Plant Physiol. Biochem.* **49**, 782–791.
- Feys, B., Benedetti, C.E., Penfold, C.N. and Turner, J.G. (1994) *Arabidopsis* mutants selected for resistance to the phytotoxin coronatine are male sterile, insensitive to methyl jasmonate, and resistant to a bacterial pathogen. *Plant Cell*, **6**, 751–759.
- Fonseca, S., Chini, A., Hamburger, M., Adie, B., Porzel, A., Kramell, R., Miersch, O., Wasternack, C. and Solano, R. (2009) (+)-7-*iso*-Jasmonoyl-L-isoleucine is the endogenous bioactive jasmonate. *Nat. Chem. Biol.* **5**, 344–350.
- Glauser, G., Dubugnon, L., Mousavi, S.A., Rudaz, S., Wolfender, J.L. and Farmer, E.E. (2009) Velocity estimates for signal propagation leading to systemic jasmonic acid accumulation in wounded *Arabidopsis*. *J. Biol. Chem.* **284**, 34506–34513.
- Gutierrez, L., Mongelard, G., Flokova, K. et al. (2012) Auxin controls *Arabidopsis* adventitious root initiation by regulating jasmonic acid homeostasis. *Plant Cell*, **24**, 2515–2527.
- Gutjahr, C., Riemann, M., Müller, A., Düchting, P., Weiler, E.W. and Nick, P. (2005) Cholodny-Went revisited: a role for jasmonate in gravitropism of rice coleoptiles. *Planta*, **222**, 575–585.
- Hoffmann, M., Hentrich, M. and Pollmann, S. (2011) Auxin-oxylipin cross-talk: relationship of antagonists. *J. Integr. Plant Biol.* **53**, 429–445.
- Howe, G.A. and Jander, G. (2008) Plant immunity to insect herbivores. *Annu. Rev. Plant Biol.* **59**, 41–66.
- Hruz, T., Laule, O., Szabo, G., Wessendorp, F., Bleuler, S., Oertle, L., Widmayer, P., Gruissem, W. and Zimmermann, P. (2008) Genevestigator V3: a reference expression database for the meta-analysis of transcriptomes. *Adv. Bioinformatics*. 420747
- Hull, A.K., Vij, R. and Celenza, J.L. (2000) *Arabidopsis* cytochrome P450s that catalyze the first step of tryptophan-dependent indole-3-acetic acid biosynthesis. *Proc. Natl Acad. Sci. USA*, **97**, 2379–2384.
- Ishiguro, S., Kawai-Oda, A., Ueda, J., Nishida, I. and Okada, K. (2001) The *DEFECTIVE IN ANTHEP DEHISCENCE1* gene encodes a novel phospholipase A1 catalyzing the initial step of jasmonic acid biosynthesis, which synchronizes pollen maturation, anther dehiscence, and flower opening in *Arabidopsis*. *Plant Cell*, **13**, 2191–2209.
- Kawashima, T. and Goldberg, R.B. (2010) The suspensor: not just suspending the embryo. *Trends Plant Sci.* **15**, 23–30.
- Kazan, K. and Manners, J.M. (2008) Jasmonate signaling: toward an integrated view. *Plant Physiol.* **146**, 1459–1468.
- Kim, J.I., Sharkhuu, A., Jin, J.B. et al. (2007) *yucca6*, a dominant mutation in *Arabidopsis*, affects auxin accumulation and auxin-related phenotypes. *Plant Physiol.* **145**, 722–735.
- Kim, J.I., Murphy, A.S., Baek, D., Lee, S.W., Yun, D.J., Bressan, R.A. and Narasimhan, M.L. (2011) *YUCCA6* over-expression demonstrates auxin function in delaying leaf senescence in *Arabidopsis thaliana*. *J. Exp. Bot.* **62**, 3981–3992.
- King, J.J., Stimart, D.P., Fisher, R.H. and Bleeker, A.B. (1995) A mutation altering auxin homeostasis and plant morphology in *Arabidopsis*. *Plant Cell*, **7**, 2023–2037.
- Kubigsteltig, I., Laudert, D. and Weiler, E.W. (1999) Structure and regulation of the *Arabidopsis thaliana* allene oxide synthase gene. *Planta*, **208**, 463–471.
- LeClere, S., Schmelz, E.A. and Chourey, P.S. (2010) Sugar levels regulate tryptophan-dependent auxin biosynthesis in developing maize kernels. *Plant Physiol.* **153**, 306–318.
- Lehmann, T. and Pollmann, S. (2009) Gene expression and characterization of a stress-induced tyrosine decarboxylase from *Arabidopsis thaliana*. *FEBS Lett.* **583**, 1895–1900.
- Lehmann, T., Hoffmann, M., Hentrich, M. and Pollmann, S. (2010) Indole-3-acetamide-dependent auxin biosynthesis: a widely distributed way of indole-3-acetic acid production? *Eur. J. Cell Biol.* **89**, 895–905.
- Liu, C.M. and Meinke, D.W. (1998) The *titan* mutants of *Arabidopsis* are disrupted in mitosis and cell cycle control during seed development. *Plant J.* **16**, 21–31.
- Livak, K.J. and Schmittgen, T.D. (2001) Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta CT}$  method. *Methods*, **25**, 402–408.
- Mandaokar, A., Thines, B., Shin, B., Lange, B.M., Choi, G., Koo, Y.J., Yoo, Y.J., Choi, Y.D. and Browse, J. (2006) Transcriptional regulators of stamen development in *Arabidopsis* identified by transcriptional profiling. *Plant J.* **46**, 984–1008.
- Mashiguchi, K., Tanaka, K., Sakai, T. et al. (2011) The main auxin biosynthesis pathway in *Arabidopsis*. *Proc. Natl Acad. Sci. USA*, **108**, 18512–18517.
- Mueller, S., Hilbert, B., Dueckershoff, K., Roitsch, T., Kruschke, M., Mueller, M.J. and Berger, S. (2008) General detoxification and stress responses are mediated by oxidized lipids through TGA transcription factors in *Arabidopsis*. *Plant Cell*, **20**, 768–785.
- Nagpal, P., Ellis, C.M., Weber, H. et al. (2005) Auxin response factors ARF6 and ARF8 promote jasmonic acid production and flower maturation. *Development*, **132**, 4107–4118.

- Novák, O., Hényková, E., Sairanen, I., Kowalczyk, M., Pospíšil, T. and Ljung, K. (2012) Tissue-specific profiling of the *Arabidopsis thaliana* auxin metabolome. *Plant J.* **72**, 523–536.
- Pauwels, L., Morreel, K., De Witte, E., Lammertyn, F., Van Montagu, M., Boerjan, W., Inze, D. and Goossens, A. (2008) Mapping methyl jasmonate-mediated transcriptional reprogramming of metabolism and cell cycle progression in cultured *Arabidopsis* cells. *Proc. Natl Acad. Sci. USA*, **105**, 1380–1385.
- Pauwels, L., Barbero, G.F., Geerinck, J. *et al.* (2010) NINJA connects the co-repressor TOPLESS to jasmonate signalling. *Nature*, **464**, 788–791.
- Pollmann, S., Neu, D., Lehmann, T., Berkowitz, O., Schäfer, T. and Weiler, E.W. (2006) Subcellular localization and tissue specific expression of amidase 1 from *Arabidopsis thaliana*. *Planta*, **224**, 1241–1253.
- Pollmann, S., Dücking, P. and Weiler, E.W. (2009) Tryptophan-dependent indole-3-acetic acid biosynthesis by 'IAA-synthase' proceeds via indole-3-acetamide. *Phytochemistry*, **70**, 523–531.
- Ramakers, C., Ruijter, J.M., Deprez, R.H. and Moorman, A.F. (2003) Assumption-free analysis of quantitative real-time polymerase chain reaction (PCR) data. *Neurosci. Lett.* **339**, 62–66.
- Rampey, R.A., LeClere, S., Kowalczyk, M., Ljung, K., Sandberg, G. and Bartel, B. (2004) A family of auxin-conjugate hydrolases that contributes to free indole-3-acetic acid levels during *Arabidopsis* germination. *Plant Physiol.* **135**, 978–988.
- Rawat, R., Schwartz, J., Jones, M.A., Sairanen, I., Cheng, Y., Andersson, C.R., Zhao, Y., Ljung, K. and Harmer, S.L. (2009) REVEILLE1, a Myb-like transcription factor, integrates the circadian clock and auxin pathways. *Proc. Natl Acad. Sci. USA*, **106**, 16883–16888.
- Ren, C., Pan, J., Peng, W. *et al.* (2005) Point mutations in *Arabidopsis Cullin1* reveal its essential role in jasmonate response. *Plant J.* **42**, 514–524.
- Riemann, M., Müller, A., Korte, A., Furuya, M., Weiler, E.W. and Nick, P. (2003) Impaired induction of the jasmonate pathway in the rice mutant *hebiba*. *Plant Physiol.* **133**, 1820–1830.
- Staswick, P.E., Su, W. and Howell, S.H. (1992) Methyl jasmonate inhibition of root growth and induction of a leaf protein are decreased in an *Arabidopsis thaliana* mutant. *Proc. Natl Acad. Sci. USA*, **89**, 6837–6840.
- Stavang, J.A., Gallego-Bartolomé, J., Gómez, M.D., Yoshida, S., Asami, T., Olsen, J.E., García-Martínez, J.L., Alabadi, D. and Blázquez, M.A. (2009) Hormonal regulation of temperature-induced growth in *Arabidopsis*. *Plant J.* **60**, 589–601.
- Stepanova, A.N., Yun, J., Robles, L.M., Novak, O., He, W., Guo, H., Ljung, K. and Alonso, J.M. (2011) The *Arabidopsis* YUCCA1 flavin monooxygenase functions in the indole-3-pyruvic acid branch of auxin biosynthesis. *Plant Cell*, **23**, 3961–3973.
- Sugawara, S., Hishiyama, S., Jikumaru, Y., Hanada, A., Nishimura, T., Koshida, T., Zhao, Y., Kamiya, Y. and Kasahara, H. (2009) Biochemical analyses of indole-3-acetaldoxime-dependent auxin biosynthesis in *Arabidopsis*. *Proc. Natl Acad. Sci. USA*, **106**, 5430–5435.
- Sun, J., Xu, Y., Ye, S. *et al.* (2009) *Arabidopsis* ASA1 is important for jasmonate-mediated regulation of auxin biosynthesis and transport during lateral root formation. *Plant Cell*, **21**, 1495–1511.
- Sun, J., Chen, Q., Qi, L. *et al.* (2011) Jasmonate modulates endocytosis and plasma membrane accumulation of the *Arabidopsis* PIN2 protein. *New Phytol.* **191**, 360–375.
- Szemenyei, H., Hannon, M. and Long, J.A. (2008) TOPLESS mediates auxin-dependent transcriptional repression during *Arabidopsis* embryogenesis. *Science*, **319**, 1384–1386.
- Tao, Y., Ferrer, J.L., Ljung, K. *et al.* (2008) Rapid synthesis of auxin via a new tryptophan-dependent pathway is required for shade avoidance in plants. *Cell*, **133**, 164–176.
- Thimann, K.V. (1938) Hormones and the analysis of growth. *Plant Physiol.* **13**, 437–449.
- Tiryaki, I. and Staswick, P.E. (2002) An *Arabidopsis* mutant defective in jasmonate response is allelic to the auxin-signaling mutant *axr1*. *Plant Physiol.* **130**, 887–894.
- Tobeña-Santamaría, R., Bliet, M., Ljung, K., Sandberg, G., Mol, J.N., Souer, E. and Koes, R. (2002) FLOOZY of petunia is a flavin mono-oxygenase-like protein required for the specification of leaf and flower architecture. *Genes Dev.* **16**, 753–763.
- Truman, W.M., Bennett, M.H., Turnbull, C.G. and Grant, M.R. (2010) *Arabidopsis* auxin mutants are compromised in systemic acquired resistance and exhibit aberrant accumulation of various indolic compounds. *Plant Physiol.* **152**, 1562–1573.
- Uppalapati, S.R., Ayoubi, P., Weng, H., Palmer, D.A., Mitchell, R.E., Jones, W. and Bender, C.L. (2005) The phytotoxin coronatine and methyl jasmonate impact multiple phytohormone pathways in tomato. *Plant J.* **42**, 201–217.
- Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paupe, A. and Speleman, F. (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.* **3**, RESEARCH0034.
- Wasternack, C. (2007) Jasmonates: an update on biosynthesis, signal transduction and action in plant stress response, growth and development. *Ann. Bot.* **100**, 681–697.
- Wolters, H. and Jürgens, G. (2009) Survival of the flexible: hormonal growth control and adaptation in plant development. *Nat. Rev. Genet.* **10**, 305–317.
- Won, C., Shen, X., Mashiguchi, K., Zheng, Z., Dai, X., Cheng, Y., Kasahara, H., Kamiya, Y., Chory, J. and Zhao, Y. (2011) Conversion of tryptophan to indole-3-acetic acid by TRYPTOPHAN AMINOTRANSFERASES OF *ARABIDOPSIS* and YUCCAs in *Arabidopsis*. *Proc. Natl Acad. Sci. USA*, **108**, 18518–18523.
- Woodward, C., Bemis, S.M., Hill, E.J., Sawa, S., Koshida, T. and Torii, K.U. (2005) Interaction of auxin and ERECTA in elaborating *Arabidopsis* inflorescence architecture revealed by the activation tagging of a new member of the YUCCA family putative flavin monooxygenases. *Plant Physiol.* **139**, 192–203.
- Yamamoto, Y., Kamiya, N., Morinaka, Y., Matsuoka, M. and Sazuka, T. (2007) Auxin biosynthesis by the YUCCA genes in rice. *Plant Physiol.* **143**, 1362–1371.
- Zhang, Y. and Turner, J.G. (2008) Wound-induced endogenous jasmonates stunt plant growth by inhibiting mitosis. *PLoS ONE*, **3**, e3699.
- Zhang, Z., Li, Q., Li, Z., Staswick, P.E., Wang, M., Zhu, Y. and He, Z. (2007) Dual regulation role of *GH3.5* in salicylic acid and auxin signaling during *Arabidopsis-Pseudomonas syringae* interaction. *Plant Physiol.* **145**, 450–464.
- Zhao, Y. (2008) The role of local biosynthesis of auxin and cytokinin in plant development. *Curr. Opin. Plant Biol.* **11**, 16–22.
- Zhao, Y., Christensen, S.K., Fankhauser, C., Cashman, J.R., Cohen, J.D., Weigel, D. and Chory, J. (2001) A role for flavin monooxygenase-like enzymes in auxin biosynthesis. *Science*, **291**, 306–309.
- Zhao, Y., Hull, A.K., Gupta, N.R., Goss, K.A., Alonso, J., Ecker, J.R., Normanly, J., Chory, J. and Celenza, J.L. (2002) Trp-dependent auxin biosynthesis in *Arabidopsis*: involvement of cytochrome P450s CYP79B2 and CYP79B3. *Genes Dev.* **16**, 3100–3112.